

Sleep and Health— A Lifestyle Medicine Approach

Param Dedhia, MD; Robert Maurer, PhD

doi: 10.12788/jfp.0295

INTRODUCTION

Sleep is vital for health and healing, yet it may not be getting the attention it deserves as a requirement for physical as well as mental and emotional health. Insufficient or disordered sleep is associated with serious disease, morbidity, and mortality.¹ Moreover, poor sleep has presented challenges to public health and safety. It is also the foundation upon which other lifestyle therapies, such as diet and exercise, are improved. It is very difficult for patients to adhere to a healthy diet and exercise when fatigued and not afforded mental clarity.²

The perspective of sleep as preventive medicine is furthered by appreciating its 2-way impact: Poor sleep increases the risk of disease and illness, as well as the converse, disease and illness disrupt sleep. This often creates a vicious cycle in which the cumulative effect is deepened morbidity and mortality.³ Modern medicine has developed treatments with a focus on pharmacology and interventions that have been helpful. Yet, for the family physician, the burden and the growth of sleep challenges will require reframing with a focus on prevention.

RISKS ASSOCIATED WITH POOR SLEEP

Sleep disorders negatively impact both short- and long-term health. The more immediate effects reduce a sense of well-being and performance.⁴ Moreover, excessive daytime sleepiness is commonly experienced, although not always recognized and/or connected to poor sleep. Accumulated effects

Param Dedhia, MD¹

Robert Maurer, PhD²

AUTHOR AFFILIATIONS

¹ MOVEO Health, Tucson, AZ

² Family Medicine Spokane, Spokane, WA

DISCLOSURES

The authors have no conflicts of interest to disclose.

of disordered sleep include premature mortality, cardiovascular disease, hypertension, obesity, metabolic syndrome, diabetes and impaired glucose tolerance, immunosuppression, inflammation, cancer, cognitive impairment, and psychiatric disorders such as anxiety and depression.⁵

SLEEP AND OBESITY

Today, we are witnessing 2 epidemics: increasing obesity and increasing sleep disorders.^{6,7} Obesity is reaching overwhelming proportions throughout the developed world and is attributed largely to industrialization, increased food consumption, and lower levels of physical activity.^{7,8} The role of sleep in obesity is becoming increasingly understood. Sleep deprivation and disorders have been hypothesized to contribute toward obesity by decreasing leptin, increasing ghrelin, and compromising insulin sensitivity.⁹ There is a negative relationship between sleep duration and central adiposity. This has been recognized as a significant risk factor in the pathophysiology of obstructive sleep apnea in adults. Furthermore, obstructive sleep apnea is associated with increased body mass index.¹⁰

SLEEP AND HEART DISEASE

Atherosclerotic cardiovascular disease (ASCVD) is one of the most prevalent diseases in industrial nations. Even with an improved ability to diagnose and treat ASCVD, the disease and its consequences are important contributors to morbidity and mortality. Therefore, it is necessary to go beyond the management of traditional ASCVD risk factors and seek other factors and comorbidities that might contribute to its development and progression.¹¹

SLEEP AND DIABETES

The increasing prevalence of type 2 diabetes (T2D) can be attributed to dramatic lifestyle changes in response to the industrialization of modern society that may not be limited to changes in diet and physical activity.¹² As with cardiovascular disease, one such factor strongly associated with the

development and progression of T2D is sleep. Population studies have observed a U-shaped relationship between sleep duration and T2D risk; those who self-report habitually sleeping less than 7 hours or more than 8 hours are at increased risk.¹³ Decreased insulin sensitivity due to short sleep duration is observed among patients and in laboratory studies.¹⁴⁻²³ Furthermore, when sleep time is extended in short sleepers, insulin sensitivity improves.²⁴

SLEEP AND IMMUNITY/INFLAMMATION

Poor immune status and increased inflammation are also associated with poor quantity or quality of sleep. There are no clear studies indicating whether inflammation causes poor sleep or the reverse. However, the combination of poor immune status and increased inflammation puts patients at risk for poor sleep and poor health. It is appropriate for the immune system to be turned on in the setting of infection or illness, but inflammation may be observed when the immune system is triggered. It is increasingly appreciated that lifestyle practices, especially poor sleep, directly impact both inflammation and immunocompetence.²⁵⁻²⁷

As vaccinations have been a cornerstone of preventive medicine, it is important to draw the connections between sleep and vaccinations. Sleep promotes antiviral immunity by supporting the adaptive immune response,²⁸ with evidence that experimental and naturalistic sleep loss is associated with poorer immunologic memory after a vaccination.²⁹⁻³¹ For example, one may not achieve the full benefit of the hepatitis B series as well as the hepatitis A and influenza vaccinations if followed by less than 6 hours of overnight sleep.

SLEEP AND SAFETY

Sleep problems are associated with accidents and human errors.³² Insomnia and poor sleep are major contributors to unintentional fatal injuries in general as well as in fatal motor vehicle injuries.³³ Traffic accidents and injuries among shift workers are also more likely to occur during nighttime hours. This surges around 2:00 to 3:00 AM, when there is the greatest tendency toward sleep with the circadian rhythm.³⁴

BENEFITS OF HEALTHY SLEEP

The casual view of sleep as simply a dormant and passive unconsciousness with the suspension of normal bodily activities shifted as neurology laid the foundation for understanding sleep using electroencephalography (EEG). The brain is very active during sleep, in which vital restoration of the mind and body occurs with each night's rest. Sleep affects our daily functioning and is essential to our physical, mental, and emotional health. William Shakespeare so insightfully

and aptly described sleep as "nature's soft nurse." Quality sleep improves learning, memory, and mood and enhances motivation for other lifestyle-enhancing behaviors, such as exercise and healthy food choices.³⁵⁻³⁸

AMOUNT OF SLEEP NEEDED

The simple response to the question of "how much sleep do I need?" is the sleep time that permits a person to be wide awake, alert, and energetic throughout the day without the aid of stimulants such as coffee. The vast amount of the adult population requires about 8 hours of sleep.

The National Sleep Foundation Scientific Advisory Council has recommended sleep ranges for all age groups (see **TABLE 1**).³⁹

Optimal sleep for an individual varies from person to person and during their lifetime. Moreover, some adults do not fit into the guidelines for optimal sleep. Requiring more than 9 hours of sleep (being a "long sleeper") or needing less than 6 hours (being a "short sleeper") does not reflexively diagnose an individual with a sleep disorder. There are genetic predispositions that allow people to be outside of the recommended sleep parameters and have normal and healthy daytime functioning. Approximately 5% to 10% of the adult population are "long sleepers," and about 5% function well as "short sleepers."³⁹

ASSESSING SLEEP CHALLENGES

Although more than half of primary care patients may experience insomnia, only about one-third report this problem to their physicians. With only 5% of people seeking treatment,^{40,41} the vast majority of people with insomnia remain untreated.⁴² Given the fast pace of primary care visits and the time needed to understand underlying etiology, it is not surprising that two-thirds of patients with insomnia report a poor understanding of treatment options, and many turn to alcohol (28%) or untested over-the-counter remedies (23%).⁴⁰

Asking patients about daytime fatigue is likely to elicit reports of sleep problems. In addition to daytime fatigue, the presenting problems may include anxiety, depression, loss of libido, hypertension, lack of concentration, concerns about possible attention-deficit/hyperactivity disorder, weight gain, relationship problems, and concerns about memory loss. Before initiating pharmacologic and/or behavioral treatment, it is important to rule out a few common and often overlooked etiologies for poor sleep. These include (1) circadian rhythm disorders, (2) eating habits, and (3) poor sleep hygiene.

A brief interview is often sufficient to assess for circadian rhythm disorders. When asking the patient, "Do you consider yourself a night owl?" or "If you did not have early morning

responsibilities, when would you prefer to go to sleep?" you are listening for those who prefer early bedtimes or those who prefer to go to bed at midnight or later. It is the mismatch between the body's preferred bedtime and scheduling demands that is causing the sleep problem.

Eating patterns and food choices influence overall health as well as sleep health. Individuals consuming an excessive number of calories report short sleep time and quality.⁴³ Concentrated carbohydrates such as sugars, just like caffeine, act as stimulants on the body, influencing a wide range of neurotransmitter shifting that makes the ability to fall asleep and stay asleep more difficult.⁴⁴ Individual variance in food tolerance, such as spicy foods and dairy, also impacts the ability to physically be soothed to be able to sleep. Large meals eaten close to bedtime typically disrupt sleep onset and/or sleep quality. As discussed earlier, poor sleep creates the hormonal and neurochemical basis for food cravings. Again, we see the vicious cycle of poor sleep leading to both overconsumption and poor food choices, limiting restorative sleep.

Sleep hygiene issues such as depriving oneself of sleep to enjoy nighttime activities and the use of electronics late into the night can create sleep difficulties that patients may be willing to modify.

TREATMENT OF INSOMNIA

The paradigm of therapy starts with etiology: comorbid insomnia due to another sleep disorder or a medical disorder that requires treatment of the underlying process or the more common psychophysiological insomnia requiring cognitive and behavioral approaches. Cognitive behavioral therapy for insomnia (CBT-I), which is a well-established, evidence-based, and efficacious treatment for insomnia,⁴⁵⁻⁴⁸ is commonly prescribed for depression. However, clinical trials have shown it is the most effective long-term solution for those with insomnia.⁴⁹ Patients already on a prescribed sleep aid can be tapered off the drug and started on CBT-I concurrently.

The positive effects of CBT-I on sleep quality are robust over time.^{50,51} CBT-I has been found to be 70%-80% efficacious in populations with a variety of comorbid medical conditions,⁵² including comorbid insomnia,⁵³ comorbid psychiatric conditions,⁵⁴ and chronic pain.⁵⁵⁻⁵⁸

CBT-I helps identify the negative attitudes and beliefs that hinder sleep and replaces them with positive thoughts, effectively "unlearning" the negative beliefs.⁵⁹ The behavioral aspect of CBT-I focuses on helpful sleep habits and avoiding unhelpful sleep behaviors. Behavioral techniques—CBT-I over a period of 6-8 weekly sessions for most adults in either individualized- or group-based administration of CBT-I—have been shown to be effective,^{52,60,64} yet these techniques are greatly underutilized in comparison to pharmacologic

TABLE 1. Recommended sleep duration by age group³⁹

Age group	Sleep hours per day
Newborns (0-3 months)	14-17
Infants (4-11 months)	12-15
Toddlers (1-2 years)	11-14
Preschoolers (3-5 years)	10-13
School-age children (6-13 years)	9-11
Teenagers (14-17)	8-10
Younger adults (18-25)	7-9
Adults (26-64)	7-9
Older adults (≥65)	7-8

approaches. There is an app called CBT-I Coach that is both evidence-based and available at no cost.⁶⁵ More recently, digital cognitive behavioral therapy for insomnia was shown to promote later health resilience during the coronavirus pandemic.⁶⁵

LIFESTYLE AS TREATMENT

The important impact of lifestyle behaviors on sleep must be appreciated. Supporting the patient's connection to their environment, healthy nutrition, exercise, and stress management provides opportunities for better health and sleep.

The impact of the environment on sleep health is highlighted by the effect of the diurnal light and darkness cycle on sleep quality and duration. Light is the strongest synchronizing agent for the circadian system. Moreover, it is the strongest external cue to stimulate the reticular activating system in the brain and alertness. A proposed mechanism includes the suppression of endogenous melatonin. Blood levels of the pineal hormone melatonin are high at night and low during the day.⁶⁶ A cornerstone of healthy sleep is routine, as well as regular patterns. The modern era, with digital screens and 24-hour expectations, has challenged our physiology to promote sleep. As melatonin production is inherently reduced from adolescence to adulthood, this begins to explain why some individuals benefit from supplementation of melatonin to induce and promote sleep. For this to be effective, partnership and buy-in from the patient are essential given the commitment needed. Furthermore, the family physician may wish to collaborate with a sleep medicine specialist, given the complexity of dosing and timing.

In recent years, many nutritional supplements have been used to benefit sleep wellness. However, the relationship between nutritional components and sleep is complicated. Nutritional factors vary dramatically with different

diet patterns and depend significantly on the digestive and metabolic functions of each individual. Moreover, nutrition can significantly affect the hormones and inflammation status that directly or indirectly contribute to insomnia. With the rise of personalized medicine and personalized nutrition, there has been a growing body of research and clinical experience on individualizing nutritional factors, carbohydrates, lipids, amino acids, and vitamins to promote sleep and reduce sleep disorders.⁶⁷ Simply put, nutrition and dietetics are important opportunities for better sleep health.

The National Sleep Foundation's 2013 Sleep in America poll highlighted the association between exercise and better sleep.⁶⁸ It is thought that a physically active daytime uses adenosine triphosphate resources such that the cleaving of the phosphate bonds results in a higher amount of adenosine by bedtime. Adenosine promotes sleep induction and deep sleep stages. Exercisers, compared with nonexercisers, are more likely to report restorative sleep. Poor sleep makes us less likely to exercise, which in turn leads to relative difficulty falling asleep or falling back asleep in the middle of the night and waking up too early.⁶⁹⁻⁷¹ Thus, there is a vicious cycle of reduced physical activity and reduced sleep. Although the timing of exercise has been widely debated, it is likely to be based on individual experience. Regardless, daily physical activity promotes nightly rest.

Stress and sleep are closely related as a result of the substantial overlap in neurotransmitter signaling and regulatory pathways between the neural centers that modulate mood and the sleep-wake cycle. Both acute and chronic stressors, and individual variability in coping with stress, are major determinants of sleep quality and quantity. Different approaches to stress reduction demonstrate opportunities to promote sleep onset, sleep maintenance, and daytime robustness. As with both nutrition and exercise, there are different levels of sleep benefits with stress reduction. Individualizing approaches offers a greater likelihood for sleep health and overall health.^{72,73}

As outlined in this article, there is a great opportunity to help patients see the relationship between successful sleep and their food intake, exercise, and stress management. As this is the province of preventive lifestyle medicine, it is a key to optimal health. Some patients approach sleep using pharmaceutical aids, and there is an opportunity for family physicians to educate and offer patients resources for healthy sleep. Lifestyle medicine and healthy sleep are essential pillars that we can offer to all of our patients for true health and healing. ●

REFERENCES

1. Cacho V, Lum E. *Integrative Sleep Medicine*. Oxford, UK: Oxford University Press; 2021.
2. Walker M. *Why We Sleep*. London, UK: Penguin Books; 2018.
3. Worley S. The extraordinary importance of sleep: the detrimental effects of inadequate sleep on health and public safety drive an explosion of sleep research. *P T*. 2018;43(12):758-763.
4. Medic G, Wille M, Hemels M. Short- and long-term health consequences of sleep disruption. *Nat Sci Sleep*. 2017;9:151-161.
5. Institute of Medicine, Board on Health Sciences Policy, Committee on Sleep Medicine and Research. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. Washington, DC: National Academies Press; 2006.
6. Adult Sleep Habits. National Sleep Foundation. Published 2021. Accessed August 17, 2021. <https://www.sleepfoundation.org/professionals/sleep-american-polls/2002-adult-sleep-habits>
7. Stein C, Colditz G. The epidemic of obesity. *J Clin Endocrinol Metab*. 2004;89(6):2522-2525. doi:10.1210/jc.2004-0288
8. Eskin SB, Hermanson S. Nutrition labeling at fast-food and other chain restaurants. *Issue Brief (Public Policy Inst (Am Assoc Retired Pers))*. 2004;(1b71):1-6.
9. Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB. Inadequate sleep as a risk factor for obesity: analyses of the NHANES I. *Sleep*. 2005;28(10):1289-1296. doi:10.1093/sleep/28.10.1289
10. Bonsignore MR, McNicholas WT, Montserrat JM, Eckel J. Adipose tissue in obesity and obstructive sleep apnoea. *Eur Respir J*. 2012;39(3):746-767. doi:10.1183/09031936.00047010
11. Arzt M, Hetzencker A, Steiner S, Buchner S. Sleep-disordered breathing and coronary artery disease. *Can J Cardiol*. 2015;31(7):909-917. doi:10.1016/j.cjca.2015.03.032
12. Qian J, Scheer E. Circadian system and glucose metabolism: implications for physiology and disease. *Trends Endocrinol Metab*. 2016;27(5):282-293. doi:10.1016/j.tem.2016.03.005
13. Shan Z, Ma H, Xie M, et al. Sleep duration and risk of type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care*. 2015;38(3):529-537. doi:10.2337/dc14-2073
14. Buxton OM, Pavlova M, Rei EW, Wang W, Simonson DC, Adler GK. Sleep restriction for 1 week reduces insulin sensitivity in healthy men. *Diabetes*. 2010;59(9):2126-2133. doi:10.2337/db09-0699
15. Darukhanavala A, Booth JN 3rd, Bromley L, Whitmore H, Imperial J, Penev PD. Changes in insulin secretion and action in adults with familial risk for type 2 diabetes who curtail their sleep. *Diabetes Care*. 2011;34(10):2259-2264. doi:10.2337/dc11-0777
16. Donga E, van Dijk M, van Dijk JG, et al. Partial sleep restriction decreases insulin sensitivity in type 1 diabetes. *Diabetes Care*. 2010;33(7):1573-1577. doi:10.2337/dc09-2317
17. Donga E, van Dijk M, van Dijk JG, et al. A single night of partial sleep deprivation induces insulin resistance in multiple metabolic pathways in healthy subjects. *J Clin Endocrinol Metab*. 2010;95(6):2963-2968. doi:10.1210/jc.2009-2430
18. Klingenberg L, Chaput JP, Holmbäck U, et al. Acute sleep restriction reduces insulin sensitivity in adolescent boys. *Sleep*. 2013;36(7):1085-1090. doi:10.5665/sleep.2816
19. Matthews KA, Dahl RE, Owens JF, Lee L, Hall M. Sleep duration and insulin resistance in healthy black and white adolescents. *Sleep*. 2012;35(10):1353-1358. doi:10.5665/sleep.2112
20. Nedeltcheva AV, Kessler L, Imperia J, Penev PD. Exposure to recurrent sleep restriction in the setting of high caloric intake and physical inactivity results in increased insulin resistance and reduced glucose tolerance. *J Clin Endocrinol Metab*. 2009;94(9):3242-3250. doi:10.1210/jc.2009-0483
21. Robertson MD, Russell-Jone D, Umpley AM, Dijk DJ. Effects of three weeks of mild sleep restriction implemented in the home environment on multiple metabolic and endocrine markers in healthy young men. *Metabolism*. 2013;62(2):204-211. doi:10.1016/j.metabol.2012.07.016
22. Schmid SM, Hallschmid M, Jauch-Chara K, et al. Disturbed glucoregulatory response to food intake after moderate sleep restriction. *Sleep*. 2011;34(3):371-377. doi:10.1093/sleep/34.3.371
23. Wong PM, Manuck SB, DiNardo MM, Korytkowski M, Muldoon MF. Shorter sleep duration is associated with decreased insulin sensitivity in healthy white men. *Sleep*. 2015;38(2):223-231. doi:10.5665/sleep.4402
24. Leproult R, Deliens G, Gilson M, Peigneux P. Beneficial impact of sleep extension on fasting insulin sensitivity in adults with habitual sleep restriction. *Sleep*. 2015;38(5):707-715. doi:10.5665/sleep.4660
25. Besedovsky L, Lange T, Born J. Sleep and immune function. *Pflugers Arch*. 2012;463(1):121-137. doi:10.1007/s00424-011-1044-0
26. Dinges DF, Douglas SD, Hamarman S, Zaugg L, Kapoor S. Sleep deprivation and human immune function. *Adv Neuroimmunol*. 1995;5(2):97-110. doi:10.1016/0960-5428(95)00002-j
27. Palma BD, Gabriel A Jr, Colugnati FA, Tufik S. Effects of sleep deprivation on the development of autoimmune disease in an experimental model of systemic lupus erythematosus. *Am J Physiol Regul Integr Comp Physiol*. 2006;291(5):1527-1532. doi:10.1152/ajpregu.00186.2006
28. Gangwisch JE, Heymsfield SB, Boden-Albala B, et al. Sleep duration associated with mortality in elderly, but not middle-aged, adults in a large US sample. *Sleep*. 2008;31(8):1087-1096.
29. Lange T, Dimitrov S, Born J. Effects of sleep and circadian rhythm on the human immune system. *Ann N Y Acad Sci*. 2010;1193:48-59. doi:10.1111/j.1749-6632.2009.05300.x
30. Lange T, Perras B, Fehm HL, Born J. Sleep enhances the human antibody response to hepatitis A vaccination. *Psychosom Med*. 2003;65(5):831-835. doi:10.1097/01.psy.0000091382.61178.f1
31. Prather AA, Hall M, Furry JM, et al. Sleep and antibody response to hepatitis B vaccination. *Sleep*. 2012;35(8):1063-1069. doi:10.5665/sleep.1990

32. Akerstedt T, Fredlund P, Gillberg M, Jansson B. A prospective study of fatal occupational accidents—relationship to sleeping difficulties and occupational factors. *J Sleep Res*. 2002;11(1):69-71. doi:10.1046/j.1365-2869.2002.00287.x
33. Laugsand LE, Strand LB, Vatten LJ, Janszky I, Bjørngaard JH. Insomnia symptoms and risk for unintentional fatal injuries—the HUNT Study. *Sleep*. 2014;37(11):1777-1786. doi:10.5665/sleep.4170
34. Boivin DB, Boudreau P. Impacts of shift work on sleep and circadian rhythms. *Pathol Biol (Paris)*. 2014;62(5):292-301. doi:10.1016/j.patbio.2014.08.001
35. Dolezal BA, Neufeld EV, Boland DM, Martin JL, Cooper CB. Interrelationship between sleep and exercise: a systematic review. *Adv Prev Med*. 2017;2017:1364-1387. doi:10.1155/2017/1364387
36. Greer SM, Goldstein AN, Walker MP. The impact of sleep deprivation on food desire in the human brain. *Nat Commun*. 2013;4:2259. doi:10.1038/ncomms3259
37. Rasch B, Born J. About sleep's role in memory. *Physiol Rev*. 2013;93(2):681-766. doi:10.1152/physrev.00032.2012
38. Triantafyllou S, Saeb S, Lattie EG, Mohr DC, Kording KP. Relationship between sleep quality and mood: ecological momentary assessment study. *JMIR Ment Health*. 2019;6(3):12613. doi:10.2196/12613
39. Watson NF. Sleep duration: a consensus conference. *J Clin Sleep Med*. 2015;11(1):7-8. doi:10.5664/jcsm.4382
40. Ancoli-Israel S, Roth T. Characteristics of insomnia in the United States: results of the 1991 National Sleep Foundation Survey. I. *Sleep*. 1999;22 Suppl 2:S347-S353.
41. Shochat T, Umphress J, Israel AG, Ancoli-Israel S. Insomnia in primary care patients. *Sleep*. 1999;22 Suppl 2:S359-S365.
42. Mellinger GD, Balter MB, Uhlenhuth EH. Insomnia and its treatment. Prevalence and correlates. *Arch Gen Psychiatry*. 1985;42(3):225-232. doi:10.1001/archpsyc.1985.01790260019002
43. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Dietary nutrients associated with short and long sleep duration. Data from a nationally representative sample. *Appetite*. 2013;64:71-80. doi:10.1016/j.appet.2013.01.004
44. Beebe DW, Simon S, Sumner S, Hemmer S, Strotman D, Dolan LM. Dietary intake following experimentally restricted sleep in adolescents. *Sleep*. 2013;36(6):827-834. doi:10.5665/sleep.2704
45. Espie CA. "Stepped care": a health technology solution for delivering cognitive behavioral therapy as a first line insomnia treatment. *Sleep*. 2009;32(12):1549-1558. doi:10.1093/sleep/32.12.1549
46. McCurry SM, Logsdon RG, Teri L, Vitiello MV. Evidence-based psychological treatments for insomnia in older adults. *Psychol Aging*. 2007;22(1):18-27. doi:10.1037/0882-7974.22.1.18
47. Morin CM, Bootzin RR, Buysse DJ, Edinger JD, Espie CA, Lichstein KL. Psychological and behavioral treatment of insomnia: update of the recent evidence (1998-2004). *Sleep*. 2006;29(11):1398-1414. doi:10.1093/sleep/29.11.1398
48. Siebern AT, Manber R. New developments in cognitive behavioral therapy as the first-line treatment of insomnia. *Psychol Res Behav Manag*. 2011;4:21-28. doi:10.2147/prbm.S10041
49. Mitchell MD, Gehrman P, Perlis M, Umscheid CA. Comparative effectiveness of cognitive behavioral therapy for insomnia: a systematic review. *BMC Fam Pract*. 2012;13:40. doi:10.1186/1471-2296-13-40
50. National Institutes of Health. National Institutes of Health State of the Science Conference Statement on Manifestations and Management of Chronic Insomnia in Adults, June 13-15, 2005. *Sleep*. 2005;28(9):1049-1057. doi:10.1093/sleep/28.9.1049
51. Morin CM, Culbert JP, Schwartz SM. Nonpharmacological interventions for insomnia: a meta-analysis of treatment efficacy. *Am J Psychiatry*. 1994;151(8):1172-1180. doi:10.1176/ajp.151.8.1172
52. Morin CM. Cognitive-behavioral approaches to the treatment of insomnia. *J Clin Psychiatry*. 2004;65 Suppl 16:33-40.
53. Wu JQ, Appleman ER, Salazar RD, Ong JC. Cognitive behavioral therapy for insomnia comorbid with psychiatric and medical conditions: a meta-analysis. *JAMA Intern Med*. 2015;175(9):1461-1472. doi:10.1001/jamainternmed.2015.3006
54. Sánchez-Ortuño MM, Edinger JD. Cognitive-behavioral therapy for the management of insomnia comorbid with mental disorders. *Curr Psychiatry Rep*. 2012;14(5):519-528. doi:10.1007/s11920-012-0312-9
55. Edinger JD, Olsen MK, Stechuchak KM, Means MK, Lineberger MD, Kirby A, et al. Cognitive behavioral therapy for patients with primary insomnia or insomnia associated predominantly with mixed psychiatric disorders: a randomized clinical trial. *Sleep*. 2009;32(4):499-510. doi:10.1093/sleep/32.4.499
56. Morgan K, Gregory P, Tomeny M, David BM, Gascoigne C. Self-help treatment for insomnia symptoms associated with chronic conditions in older adults: a randomized controlled trial. *J Am Geriatr Soc*. 2012;60(10):1803-1810. doi:10.1111/j.1532-5415.2012.04175.x
57. Rybarczyk B, Mack L, Harris JH, Stepanski E. Testing two types of self-help CBT-I for insomnia in older adults with arthritis or coronary artery disease. *Rehabil Psychol*. 2011;56(4):257-266. doi:10.1037/a0025577
58. Smith MT, Huang MI, Manber R. Cognitive behavior therapy for chronic insomnia occurring within the context of medical and psychiatric disorders. *Clin Psychol Rev*. 2005;25(5):559-592. doi:10.1016/j.cpr.2005.04.004
59. Davies DR. A multiple treatment approach to the group treatment of insomnia: a follow-up study. *Behav Cogn Psychother*. 1989;17(4):323-331.
60. Germain A, Moul DE, Franze PL, et al. Effects of a brief behavioral treatment for late-life insomnia: preliminary findings. *J Clin Sleep Med*. 2006;2(4):403-406.
61. Irwin MR, Cole JC, Nicassio PM. Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+ years of age. *Health Psychol*. 2006;25(1):3-14. doi:10.1037/0278-6133.25.1.3
62. Lovato N, Lack L, Wright H, Kennaway DJ. Evaluation of a brief treatment program of cognitive behavior therapy for insomnia in older adults. *Sleep*. 2014;37(1):117-126. doi:10.5665/sleep.3320
63. Morin CM. *Insomnia: Psychological Assessment and Management*. New York, NY: Guilford Press;1993.
64. Rybarczyk B, Lopez M, Benson R, Alsten C, Stepanski E. Efficacy of two behavioral treatment programs for comorbid geriatric insomnia. *Psychol Aging*. 2002;17(2):288-298.
65. Cheng P, Casement MD, Kalmbach DA, Castelan AC, Drake CL. Digital cognitive behavioral therapy for insomnia promotes later health resilience during the coronavirus disease 19 (COVID-19) pandemic. *Sleep*. 2021;44(4):zsa258. doi:10.1093/sleep/zsa258
66. Brown GM. Light, melatonin and the sleep-wake cycle. *J Psychiatry Neurosci*. 1994;19(5):345-353.
67. Zhao M, Tuo H, Wang S, Zhao L. The effects of dietary nutrition on sleep and sleep disorders. *Mediators Inflamm*. 2020;2020:3142874. doi:10.1155/2020/3142874
68. 2013 Exercise and Sleep. National Sleep Foundation. Accessed August 17, 2021. <https://www.sleepfoundation.org/professionals/sleep-america-polls/2013-exercise-and-sleep>
69. Guilleminault C, Clerk A, Black J, Labanowski M, Pelayo R, Claman D. Non-drug treatment trials in psychophysiological insomnia. *Arch Intern Med*. 1995;155(8):838-844.
70. Kline CE. The bidirectional relationship between exercise and sleep: implications for exercise adherence and sleep improvement. *Am J Lifestyle Med*. 2014;8(6):375-379. doi:10.1177/1559827614544437
71. Passos GS, Poyares DL, Santana MG, Tufik S, Mello MT. Is exercise an alternative treatment for chronic insomnia? *Clinics (Sao Paulo)*. 2012;67(6):653-660. doi:10.6061/clinics/2012(06)17
72. Radwan B, Yanez Touzet A, Hammami S, Chaudhury D. Prolonged exposure to social stress impairs homeostatic sleep regulation. *Front Neurosci*. 2021;15:633955. doi:10.3389/fnins.2021.633955
73. Winbush NY, Gross CR, Kreitzer M. The effects of mindfulness-based stress reduction on sleep disturbance: a systematic review. *Explore (NY)*. 2007;3(6):585-591.